

Critical regularity for elliptic equations from Littlewood-Paley theory II

Denis A. Labutin

Department of mathematics, University of California, Santa Barbara, CA 93106, USA

ABSTRACT. We establish a general theorem improving regularity of solutions of elliptic pseudodifferential equations. It allows to resolve in a unified way the regularity issue for a broad class of nonlinear elliptic equations and systems appearing in different areas of geometry and analysis.

AMS subject classification: 35J60, 58J05, 58J40

1 Introduction

The purpose of this paper is to prove a rather general local regularity theorem for elliptic equations. This theorem (Theorem 1.1) implies the full C^∞ -regularity of solutions to a large class of nonlinear elliptic problems. For example, a direct consequence of the theorem is C^∞ -smoothness of weak solutions to a full scale of nonlinear equations with conformally invariant pseudodifferential operators P_k^n introduced by Paneitz, Branson, Graham, Jenne, Mason, and Sparling [16], [1], [12]. Another consequence is the critical regularity for elliptic systems with the nonlinear structure analogous to the stationary Navier-Stokes system.

In the sequel by M we always denote a smooth paracompact manifold of dimension n , and by X an open subset of M . By $\Psi^m(M)$ we denote the class of all pseudodifferential operators on M of the order m , see section 2 for the definitions.

Theorem 1.1 *Let L , P , and Q be pseudodifferential operators from $\Psi^\alpha(M)$, $\Psi^\beta(M)$, and $\Psi^\gamma(M)$ respectively. Assume that L is elliptic of order α and that*

$$(1.1) \quad \alpha, \beta, \gamma \geq 0, \quad \alpha > \beta + \gamma.$$

Let u be a distribution on M solving

$$(1.2) \quad Lu + P(V(x)Qu) = 0 \quad \text{on } M.$$

¹E-mail address: labutin@math.ucsb.edu

Assume that $V \in L^{n/(\alpha-\beta-\gamma)}(X)$ and $u \in W^{s,p}(X)$ with

$$(1.3) \quad \alpha - \beta \geq s \geq \gamma,$$

and

$$(1.4) \quad \gamma > s - \frac{n}{p} > \alpha - \beta - n.$$

Then there exists $\varepsilon > 0$, $\varepsilon = \varepsilon(n, \alpha, \beta, \gamma, s, p)$ such that

$$(1.5) \quad u \in W^{s,p+\varepsilon}(X')$$

for any $X' \subset\subset X$.

The assumption $V \in L^{n/(\alpha-\beta-\gamma)}$ is the *critical* integrability condition. Arbitrarily small improvement of the exponent allows to derive (1.5) directly from the Calderon-Zygmund estimate. Simple examples show [14] that the integrability with any power smaller than critical is not enough to conclude (1.5) with any $\varepsilon > 0$. According to the general principle, results for linear problems of type (1.2) with rough coefficients have immediate applications to the corresponding nonlinear problems

$$Lu + f(\partial, u) = 0.$$

This paper is a continuation of our earlier work [14].

The main point of Theorem 1.1 is that the regularity improvement holds for a critical equations *without any structure assumptions* on the nonlinearity provided the nonlinearity satisfies

$$(1.6) \quad VQ(u) \in L^r_{loc} \quad \text{with} \quad \infty > r > 1.$$

As we discuss below, several nonlinear problems satisfy this condition. Main requirement (1.4) in Theorem 1.1 is nothing but (1.6). Condition (1.3) is not important and always holds in nonlinear applications.

The question of local regularity for critical elliptic equations and systems is important for different areas of geometry and analysis. Many works are dedicated to this problem for various equations. The critical equations which attracted a lot of attention include, for example, Yamabe equation, the family of Q -curvature equations, H -surfaces system, system of harmonic maps from the plane, elliptic Yang-Mills system in dimension 4, and others.

The proof Theorem 1.1 works also for ψ -do between (complex) vector bundles over M . For simplicity we do not formulate the theorem in the most general form. Below we discuss classes of nonlinear problems covered and not covered by the theorem.

Here is a typical scheme for nonlinear applications of Theorem 1.1. Consider, for example, the following equation in $\mathbf{R}^4$

$$\Delta^2 u + \partial_1^2 ((\partial_1 u)^2) = 0, \quad u \in H^2_{loc}.$$

It is critical for H^2 . Setting $V = \partial_1 u$ we arrive, due to the Sobolev embedding, at

$$\Delta^2 u + \partial_1^2 (V \partial_1 u) = 0, \quad u \in H^2_{loc}, \quad V \in L^4_{loc}.$$

Apply Theorem 1.1 to derive that $u \in W^{2,2+\varepsilon}_{loc}$. Thus we improved the integrability of $\partial_1 u$ above the critical. Now repeated applications of Calderon-Zygmund and Schauder estimates to the nonlinear problem imply that $u \in C^\infty_{loc}$.

As a direct consequence of Theorem 1.1 we recover the critical regularity of stationary Navier-Stokes system established in [26], [10]. That is, take any $u = (u_1, \dots, u_n)$, $u_j \in H^{n/4}_{loc}$, solving

$$\Delta u - \mathbf{P}(u \cdot \nabla u) = 0, \quad \nabla \cdot u = 0.$$

Here $\mathbf{P}$ is the Leray projector on the space of divergence free vector fields. Operator $\mathbf{P}$ is a homogeneous Fourier multiplier of order 0 with the singularity at the origin consistent with the scaling. Such operators enjoy the same mapping properties as ψdo from Ψ^0 , cf. [18], Ch. 6. Setting, say, $V = u$ and applying Theorem 1.1 deduce that $u \in W_{loc}^{n/4, 2+\varepsilon}$. Then the application of the standard Calderon-Zygmund and Schauder estimates gives $u \in C_{loc}^\infty$. In particular, every finite energy solution $u \in H_{loc}^1$ is smooth in dimension 4.

Another consequence of Theorem 1.1 is the full regularity for a class of nonlinear equations arising in conformal geometry. Locally these equations have the ψdo $(-\Delta)^{k/2}$, $k = 1, 2, \dots$, as the leading part. Study of these equations and related conformal concepts (for example, Branson's Q -curvature) is an active area of current research in geometry, nonlinear PDEs, mathematical physics, and spectral theory, see e.g. [3] [4], [8], [9], [11]. A corollary of Theorem 1.1 is the local regularity of weak solutions to the corresponding critical nonlinear equations. For example, any $u \in H_{loc}^{n/2}$ solving

$$\Delta^{n/2}u + \operatorname{div} \left(\left(\Delta^{3/2}u \right)^{(n-2)/3} \operatorname{grad} u \right) = 0$$

is smooth (say, in dimensions 5, 8, 11, ...). For *differential* operators this was established in [6] (for minimising solutions), [25], [7] (general case). The last two works use the same idea which is based on the unique solvability of the suitable elliptic boundary value problem. In the case of ψdo this construction does not work. This provided the initial motivation for our paper. Paper [15] contains an approach to the local regularity of (1.2) based on rewriting the equation as an integral equation in the physical space. Such approach works particularly well in the case when L , P , and Q have some structure (say, the lack of x -dependence, or possession of a radial fundamental solution, or homogeneity, ...). In the case of the second order single equations DeGiorgi-Moser tools are available [2], [24].

What critical problems are not covered by Theorem 1.1? The examples are the equation for weakly harmonic maps in the plane, and the equation for H -surfaces. The weakly harmonic maps from the plane into spheres are solutions $u \in H^1(\Omega, \mathbf{S}^N)$, $\Omega \subset \mathbf{R}^2$, of the system

$$\Delta u + |\nabla u|^2 u = 0.$$

Solutions to the H -surfaces equation are the functions $u \in H^1(\Omega, \mathbf{R}^3)$ satisfying

$$\Delta u + \partial_x u \wedge \partial_y u = 0.$$

For both equations, solutions of the class H_{loc}^1 do not satisfy the main assumption (1.4) of Theorem 1.1. For example, if we utilise the form (1.2) with $P = 1$, then the nonlinearity is only in L_{loc}^1 . If, alternatively, we utilise the divergence structure of the nonlinearity to arrive at (1.2) with $P \in \Psi^1$, then the first inequality in (1.4) is violated. Therefore, one needs to exploit more information about the nonlinearity. Presently several different proofs of the full regularity for both equations are known [20], [21], [13], [22] [5], [19]. The common element in all of them is the heavier utilisation of the div-curl structure of the nonlinearity.

2 Preliminaries

In this section we review some definitions and state some estimates needed in the proof of the theorem. We refer to monographs [17], [18], and [23] for more information.

2.1 Littlewood-Paley decomposition

Let $\{\widehat{\varphi}_j\}_{j=-\infty}^{+\infty}$ be the standard smooth partition of unity in the Littlewood-Paley theory [17], [18]. Thus $\widehat{\varphi}_j = \widehat{\varphi}(\cdot/2^j)$ is supported in, say, the ring

$$\{\xi: 2^j 3/5 \leq |\xi| \leq 2^j 5/3\} \subset (B_{2^{j+1}} \setminus B_{2^{j-1}}).$$

Let P_j denote the Littlewood-Paley projection,

$$(P_j f)^\wedge = \widehat{\varphi}_j \widehat{f}, \quad f \in \mathcal{S}'.$$

We also set

$$P_{a < \cdot < b} = \sum_{j=a+1}^{b-1} P_j.$$

It follows easily that for any j and any p , $1 \leq p \leq \infty$, the operators

$$P_j: L^p \longrightarrow L^p$$

have norms bounded uniformly over j .

Distributions with the localised Fourier transform (for example any $P_j f$) enjoy the important *Bernstein inequality*. It says that for $f \in \mathcal{S}'$ and $1 \leq p \leq q \leq \infty$

$$\|f\|_q \lesssim 2^{nj((1/p)-(1/q))} \|f\|_p \quad \text{provided} \quad \text{supp } \widehat{f} \subset B_{2^j}.$$

For $s \in \mathbf{R}^1$ and $1 < p < \infty$ the Sobolev space $W^{s,p} = W^{s,p}(\mathbf{R}^n)$ consists of all distributions with the finite norm defined by

$$\|f\|_{W^{s,p}}^p = \|P_{\leq 0} f\|_p^p + \left\| \sum_{j=1}^{\infty} (2^{2js} |P_j f|^2)^{1/2} \right\|_p^p.$$

It is well known in analysis that this norm is difficult to work with. However, in this paper we will use only the following facts about $W^{s,p}$ which follow easily from the properties of P_j . Namely,

$$\begin{aligned} f \in W^{s,p} &\implies \|P_k f\|_p \lesssim \frac{\|f\|_{W^{s,p}}}{2^{sk}} \text{ for all } k \geq 0; \\ \|P_k f\|_p &\leq \frac{N}{2^{(s+\varepsilon)k}}, \quad \varepsilon > 0, \text{ for all } k \geq 0 \implies \|f\|_{W^{s,p}} \leq C_\varepsilon N. \end{aligned}$$

The central result of the Littlewood-Paley theory implies that for $s = 0, 1, \dots$ the space $W^{s,p}$ consists of all distributions with all derivatives up to the order s lying in L^p , $1 < p < \infty$. For $p = 2$ this follows from the Plancherel isometry.

If M is a smooth paracompact manifold of dimension n , then the Sobolev spaces $W^{s,p}(M)$ (and other spaces of functions) are defined via the partition of unity. If the functions are defined on $\mathbf{R}^n$ then frequently we will not write the domain of the definition.

2.2 Pseudodifferential operators

A pseudodifferential operator (ψ do for short) A of order $m \in \mathbf{R}$ on M is a linear map

$$A: C_0^\infty(M) \longrightarrow C_{loc}^\infty(M)$$

such that in any coordinate chart A is a ψ do of order m in $\mathbf{R}^n$. This means that if we take a coordinate patch

$$U \subset M, \quad \kappa: U \rightarrow \Omega, \quad \Omega \subset \mathbf{R}^n.$$

and fix two functions $\varphi, \psi \in C_0^\infty(U)$, such that $\varphi = 1$ on $\text{supp} \psi$, then for $v \in C_0^\infty(\mathbf{R}^n)$ the map

$$(2.1) \quad v \longmapsto \left(\psi A(\varphi v \circ \kappa) \right) (\kappa^{-1} x), \quad x \in \Omega,$$

is given by the formula

$$v \longmapsto \int_{\mathbf{R}^n} e^{ix\xi} a(x, \xi) \hat{v}(\xi) d\xi,$$

where a is a symbol from S^m . The latter means that $a \in C_{loc}^\infty(\mathbf{R}^n \times \mathbf{R}^n)$ and the estimate

$$|\partial_\xi^k \partial_x^l a(x, \xi)| \leq C_{k,l} (1 + |\xi|)^{m-|k|}$$

holds for all multiindices k and l . From (2.1) it follows that $\text{supp} a \subset \kappa(\text{supp} \psi) \times \mathbf{R}^n$. By $\Psi^m(M)$ we denote the set of all ψ do on M of order (at most) m . Throughout the paper we abuse notations by denoting the operator $A_{\kappa, \varphi, \psi}$ in (2.1) by the same letter A .

We say that $A \in \Psi^m(M)$ is elliptic of order $m > 0$ if in any coordinate patch as above, for any $\Omega' \subset \subset \Omega$ there are constants $C_{1,2} > 0$ the symbol of A satisfies

$$|a(x, \xi)| \geq C_1 |\xi|^m \quad \text{if } x \in \Omega', \quad |\xi| \geq C_2.$$

Any $A \in \Psi^m(M)$, $m \in \mathbf{R}$, admits a unique extension as

$$A: \mathcal{E}'(M) \longrightarrow \mathcal{D}'(M).$$

Moreover, the central result of the Calderon-Zygmund theory can be stated as the following mapping property of ψ do:

$$A: W^{s,p}(M) \longrightarrow W_{loc}^{s-m,p}(M) \quad \text{for } s \in \mathbf{R}, \quad 1 < p < \infty.$$

2.3 Estimates for pseudodifferential operators

It is easy to see that $P_k \in \Psi^{-\infty}$. For any $A \in \Psi^m$ and any k the estimate

$$(2.2) \quad \|AP_k f\|_p \leq C_{A,p} 2^{km} \|P_{k-1 \leq \cdot \leq k+1} f\|_p, \quad f \in \mathcal{S}',$$

holds for any p , $1 \leq p \leq \infty$, with the constant $C_{A,p}$ independent of k . Estimate (2.2) is a consequence of the Calderon-Zygmund theory [18], Ch. 6.

Lemma 2.1 *For any ψ do $A \in \Psi^m$, and any P_k , $k \geq 10$ the estimate*

$$(2.3) \quad \begin{aligned} \|(P_k A - A P_k) f\|_p \leq C_{A,N} \Big(& 2^{k(m-1)} \|P_{k-5 < \cdot < k+5} f\|_p \\ & + 2^{-Nk} \|P_{\leq 0} f\| \\ & + 2^{-Nk} \sum_{j=1}^{\infty} 2^{-Nj} \|P_j f\|_p \Big), \end{aligned}$$

holds for $f \in \mathcal{S}'$, and all $N > 0$, $1 \leq p \leq \infty$. The constant $C_{A,N}$ does not depend on k .

It is easy to see, that despite the symbol $\varphi_k \in S^{-\infty}$, it admits the pointwise estimates *independent of k* only in S^0 . It is due to this fact that we can improve only by the factor 2^{-k} in (2.3).

PROOF. **1.** Let a be the symbol of A , $a \in S^m$. Let c be the symbol of $P_k A$. The basic theorem of ψ do-calculus asserts that $c \in S^{-\infty}$, and moreover

$$c(x, \xi) = \varphi(2^{-k}\xi)a(x, \xi) + \rho_k(x, \xi)$$

with $\rho_k \in S^{-\infty}$. However, we are interested in explicit dependence on k . We claim that for any multiindices α and β , $N > 0$, and all $k \geq 10$

$$(2.4) \quad \begin{aligned} |\partial_\xi^\alpha \partial_x^\beta \rho_k(x, \xi)| &\leq C_{N,A,\alpha,\beta} \\ &\times \begin{cases} (1 + |\xi|)^{m-1-|\alpha|} & \text{if } 2^{k-3} \leq |\xi| \leq 2^{k+3} \\ (1 + |\xi|)^{-N} & \text{if } |\xi| \geq 2^{k+3} \\ 2^{-Nk} & \text{if } |\xi| \leq 2^{k-3}. \end{cases} \end{aligned}$$

The point here is that $C_{N,A,\alpha,\beta}$ is independent of k .

It is obvious that estimate (2.3) follows from (2.4) and (2.2). We will prove (2.4) only for $\alpha = \beta = 0$. The obvious changes make the proof to work for the general case.

2. To show (2.4) we first write down $c(x, \xi)$. We can assume that $a(x, \xi)$ has a compact support in x . A well-known argument allows to remove this assumption while preserving (2.4), see [18], Ch. 6. We also localise P_k in x by considering

$$\varphi_{k,\varepsilon}(x, \xi) = \theta(\varepsilon x) \varphi(2^{-k}\xi),$$

where θ is a cut-off and $\varepsilon > 0$. All our estimates will be independent of ε , $0 < \varepsilon < 1$. The Fourier transform easily gives that

$$\begin{aligned} c_\varepsilon(x, \xi) &= \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} e^{ix\eta} \varphi_{k,\varepsilon}(x, \xi + \eta) \hat{a}(\eta, \xi) d\eta \\ &= \varphi_{k,\varepsilon}(x, \xi) a(x, \xi) \\ &\quad + \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} e^{ix\eta} \hat{a}(\eta, \xi) (\varphi_{k,\varepsilon}(x, \xi + \eta) - \varphi_{k,\varepsilon}(x, \xi)) d\eta, \end{aligned}$$

where $\hat{a}(\cdot, \cdot)$ denotes the Fourier transform in the *first* variable. The localisation of a in the first variable imply that

$$(2.5) \quad |\hat{a}(\eta, \xi)| \leq C_{A,N} \frac{(1 + |\xi|)^m}{(1 + |\eta|)^N}, \quad \text{for all } N.$$

Set

$$R_{k,\varepsilon}(x, \xi, \eta) = \varphi_{k,\varepsilon}(x, \xi + \eta) - \varphi_{k,\varepsilon}(x, \xi)$$

for the reminder, which we need to estimate.

3. Let us show the first estimate in (2.4). Fix any ξ satisfying $2^{k-3} \leq |\xi| \leq 2^{k+3}$. Assume first that $|\eta| \lesssim |\xi|$. Then we have

$$\begin{aligned} |R_{k,\varepsilon}(x, \xi, \eta)| &\lesssim \frac{|\eta|}{2^k} \\ &\lesssim \frac{|\eta|}{1 + |\xi|}. \end{aligned}$$

Assume next that $|\eta| \gtrsim |\xi|$. Then due to the localisation of φ_k we have

$$\begin{aligned} |R_{k,\varepsilon}(x, \xi, \eta)| &= |\varphi_{k,\varepsilon}(x, \xi)| \\ &\lesssim 1. \end{aligned}$$

Combining with (2.5) discover that

$$\begin{aligned} |\rho_{k,\varepsilon}(x, \xi)| &\lesssim C_{A,N}(1+|\xi|)^{m-1} \int_{\{|\eta| \lesssim |\xi|\}} \frac{1}{(1+|\eta|)^N} d\eta \\ &\quad + C_{A,N} \int_{\{|\eta| \gtrsim |\xi|\}} \frac{(1+|\xi|)^m}{(1+|\eta|)^N} d\eta \\ &\lesssim C_{A,N}(1+|\xi|)^{m-1}. \end{aligned}$$

Finally let $\varepsilon \rightarrow 0$. Thus the first part of (2.4) is proved.

4. Let us show the second estimate in (2.4). Fix ξ , $|\xi| \geq 2^{k+3}$. Then

$$|R_{k,\varepsilon}(x, \xi, \eta)| = 0$$

for all η such that $|\eta| \lesssim |\xi|$. Assume next that $|\eta| \gtrsim |\xi|$. Then

$$|R_{k,\varepsilon}(x, \xi, \eta)| \lesssim 1.$$

Combining with (2.5) derive that

$$\begin{aligned} |\rho_{k,\varepsilon}(x, \xi)| &\lesssim C_{A,N}(1+|\xi|)^{m-1} \int_{\{|\eta| \gtrsim |\xi|\}} \frac{1}{(1+|\eta|)^N} d\eta \\ &\lesssim C_{A,N}(1+|\xi|)^{-N'} \text{ for any } N'. \end{aligned}$$

After letting $\varepsilon \rightarrow 0$ we conclude that the second part of (2.4) is also proved.

5. To show the third estimate in (2.4) it is enough to notice that for any fixed ξ , $|\xi| \leq 2^{k-3}$, we have

$$|R_{k,\varepsilon}(x, \xi, \eta)| \neq 0 \quad \text{only for } |\eta| \asymp 2^k.$$

Hence

$$\begin{aligned} |\rho_{k,\varepsilon}(x, \xi)| &\lesssim C_{A,N} \int_{\{|\eta| \asymp 2^k\}} \frac{(1+|\xi|^m)}{(1+|\eta|)^N} d\eta \\ &\lesssim C_{A,N'} 2^{-N'k} \text{ for any } N'. \end{aligned}$$

Finally we let $\varepsilon \rightarrow 0$. ■

As a direct consequence of (2.3) we obtain that under the assumptions of Lemma 2.1 the estimate

$$(2.6) \quad \|P_k A f\|_p \leq C_{A,N} (2^{mk} \|P_{k-10 \leq \cdot \leq k+10} f\|_p + 2^{-Nk} \|f\|_p)$$

holds. Estimates (2.2) and (2.6) show that $A \in \Psi^m$ acts on the part of f with the frequencies of the order 2^k , as a multiplication by 2^{mk} modulo a fixed frequency spreading and an arbitrary small correction.

3 Proof of Theorem 1.1

By q we always denote the critical exponent,

$$q = n/(\alpha - \beta - \gamma).$$

According to (1.1) and (1.4) we have $1 < q < \infty$. In this section we write $A \lesssim B$ if $A \leq CB$ with a constant $C > 0$, depending on the operators L , P , Q , and the parameters from Theorem 1.1.

PROOF. [of Theorem 1.1] **1.** When proving (1.5) we can assume that X lies in a single coordinate chart. First we make a suitable localisation of equation (1.2). This will be done in several steps exploiting the pseudolocal character of ψdo in a standard way.

Apply definition (2.1) and utilise the pseudolocality of ψdo to derive that

$$\begin{aligned}\psi L(\varphi u) + \psi P(V(x)Qu) &= \psi L((\varphi - 1)u) \\ &= \psi f, \quad \psi f \in C_0^\infty(\Omega).\end{aligned}$$

Abusing notations we have denoted $u \circ \kappa^{-1}$ by the same letter u . Repeat this argument several times to find functions $\psi_j \in C_0^\infty(\Omega)$, $j = 1, \dots, 4$, such that $\psi_j = 1$ on $\text{supp}\psi_{j+1}$, and

$$(3.1) \quad \psi_4 L(\psi_1 u) + \psi_4 P(\psi_3 V(x)\psi_2 Q(\psi_1 u)) = \psi_4 P(\psi_3 v) + \psi_4 f,$$

where

$$v \in L^{n/(\alpha-\beta-\gamma)}(\text{supp}\psi_3), \quad f \in C_0^\infty(\text{supp}\psi_4).$$

Next, we can assume that $B_1 \subset\subset \Omega$ and

$$\psi_4 = 1 \quad \text{on} \quad B_1.$$

Take a cutoff function η_ρ ,

$$\eta_\rho = 1 \quad \text{in} \quad B_\rho, \quad \eta_\rho = 0 \quad \text{outside} \quad B_{2\rho}.$$

Later we will choose ρ small. The commutator of the multiplication by η_ρ and an operator with the symbol from S^m is a ψdo with the symbol in S^{m-1} . For differential operators this is just the Leibnitz rule. Fix a function $\phi \in C_0^\infty(\Omega)$ such that

$$\phi = 1 \quad \text{on} \quad \text{supp}\psi_1.$$

Then we can continue (3.1) and write

$$(3.2) \quad L(\eta_\rho u) + \eta_\rho P(V(x)Q(\phi u)) = \Phi_{\alpha-1}(\phi u) + \Phi_\beta v + f.$$

From now on we denote by Φ_m a generic operator from Ψ^m compactly x -supported in Ω . In (3.2) we have $f \in C_0^\infty(\Omega)$, $v \in L^q$, $\text{supp}v \subset\subset \Omega$.

Repeat this argument several times utilising $\eta_\rho = \eta_\rho \eta_{2\rho}$. Thus we discover the following relation in $\mathbf{R}^n$:

$$\begin{aligned}(3.3) \quad L(\eta_\rho u) + P(\eta_{2\rho} V(x)Q(\eta_\rho u)) &= \Phi_{\alpha-1}(\phi u) + \Phi_{\beta-1}(V(x)Q(\phi u)) \\ &+ P(\eta_{2\rho} V(x)\Phi_{\gamma-1}(\phi u)) \\ &+ \Phi_\beta v \\ &+ f,\end{aligned}$$

where the functions v and f are the same as in (3.1), (3.2).

2. Next we utilise the ellipticity to rewrite the equation in an essentially invertible form. The ellipticity of L allows us to split it,

$$L = E + M_\alpha,$$

into an operator E with the symbol $e(x, \xi) \in S^\alpha$,

$$|e(x, \xi)| \gtrsim (1 + |\xi|)^\alpha \quad \text{for all } x, \xi \in \mathbf{R}^n,$$

and the remainder M_α with the symbol $m(x, \xi) \in S^\alpha$, such that

$$\begin{aligned} m(x, \xi) &= m_1(x, \xi) + m_2(x, \xi), \\ m_1(x, \xi) &= 0, \quad \text{for } x \in B_1, \xi \in \mathbf{R}^n, \\ m_2(x, \xi) &\in S^{-\infty} \text{ is smoothing.} \end{aligned}$$

Later we will invert E modulo a smoothing ψ do via the parametrix construction. We now rewrite (3.3) as

$$\begin{aligned} E(\eta_\rho u) &= -P(\eta_{2\rho} V(x) Q(\eta_\rho u)) + \Phi_{\alpha-1}(\phi u) \\ &\quad + \Phi_{\beta-1}(V(x) Q(\phi u)) \\ &\quad + P(\eta_{2\rho} V(x) \Phi_{\gamma-1}(\phi u)) \\ &\quad + \Phi_\beta v \\ &\quad + f \\ &\quad - M_\alpha(\eta_\rho u), \end{aligned} \tag{3.4}$$

where all functions and ψ do enjoy the same properties as in (3.3).

3. We claim that

$$\eta_\rho u \in W^{\sigma, r}. \tag{3.5}$$

for some σ, r such that

$$\gamma < \sigma < \alpha - \beta \quad \text{and} \quad \sigma - \frac{n}{r} = s - \frac{n}{p}. \tag{3.6}$$

Of course, (1.3), (1.4), and (3.6) imply that $1 < r, p < \infty$. Statement (3.5) will follow rather directly from the Calderon-Zygmund estimates applied to (3.4).

Indeed, define $\nu = \alpha - \beta$. Notice that due to (1.1) $\gamma < \nu$. Rewrite (3.4) as

$$\begin{aligned} E(\eta_\rho u) &= P f_1 + \Phi_{\alpha-1} f_2 + \Phi_{\beta-1} f_3 \\ &\quad + P f_4 \\ &\quad + \Phi_\beta f_5 \\ &\quad + f_6 \end{aligned} \tag{3.7}$$

$$(3.8) \quad + M_\alpha f_7$$

Now recall the mapping properties of ψ do in $W^{s, p}$ spaces. To be able to use them we need $f_j \in L^{p_j}$ with $1 < p_j < \infty$. This is easy to verify by applying Sobolev and Holder inequalities:

$$\begin{aligned} f_1 &\in L^{p_1} \quad \text{for} \quad \frac{1}{p_1} = \frac{1}{p} + \frac{\nu}{n} - \frac{s}{n}, \\ f_2 &\in L^{p_2} \quad \text{for} \quad \frac{1}{p_2} = \frac{1}{p} - \frac{s}{n}, \\ f_3 &\in W^{p_3} \quad \text{for} \quad \frac{1}{p_3} = \frac{1}{p_1}, \\ f_4 &\in L^{p_4} \quad \text{for} \quad \frac{1}{p_4} = \frac{1}{p_1} - \frac{1}{n}, \\ f_5 &\in L^{p_5} \quad \text{for} \quad \frac{1}{p_5} = \frac{1}{q} = \frac{\nu}{n} - \frac{s}{n}. \end{aligned}$$

Our crucial assumption (1.4) ensures that all p_j are finite and $p_j \geq p_1 > 1$.

Next take a parametrix for E and apply it to both sides of (3.7). Doing so, f_6 and f_7 can be ignored because $f_6 \in C_0^\infty(\Omega)$, and in the definition of M_α we have

$$\text{supp}(\eta_\rho u) \cap \text{supp} m_1 = \emptyset.$$

The parametrix of E is a ψ do with the symbol in $S^{-\alpha}$. Utilising the mapping properties we discover that

$$\eta_\rho u = \sum_{j=1}^5 \tilde{f}_j \text{ mod } C_0^\infty(\Omega),$$

where

$$\begin{aligned} \tilde{f}_1 &\in W^{\nu, p_1} \quad \text{for} \quad \frac{1}{p_1} = \frac{1}{p} + \frac{\nu}{n} - \frac{s}{n}, \\ \tilde{f}_2 &\in W^{s+1, p_2} \quad \text{for} \quad p_2 > p_1, \\ \tilde{f}_3 &\in W^{\nu+1, p_3} \quad \text{for} \quad p_3 = p_1, \\ \tilde{f}_4 &\in W^{\nu, p_4} \quad \text{for} \quad p_4 > p_1, \\ \tilde{f}_5 &\in W^{\nu, p_5} \quad \text{for} \quad p_5 > p_1. \end{aligned}$$

Finally choose σ such that $s < \sigma < \nu$ and $\sigma < s+1$. Applying Sobolev inequalities we discover (3.5) provided

$$\frac{1}{r} = \frac{1}{p} - \frac{s}{n} + \frac{\sigma}{n}.$$

Thus (3.5), (3.6) are proved. Clearly (3.6) implies that (1.4) holds for σ and r , instead of s and p .

4. Our goal is to show that for some σ, r from (3.6) and for some $\rho > 0$ we can find $\varepsilon > 0$ such that (3.5) is improved to

$$(3.9) \quad \eta_\rho u \in W^{\sigma+\varepsilon, r}.$$

This is the main statement in the proof.

If (3.9) holds then we immediately conclude the proof of the theorem by establishing (1.5). Indeed, just repeat the arguments proving (3.5) using the improved regularity (3.9) instead of (3.6). The result will be the improved integrability of VQu . Now (1.3) allows to apply the Sobolev inequality to derive that

$$\eta_\rho u \in W^{s, p+\varepsilon'}, \quad \varepsilon' > 0.$$

The rest of the proof is devoted entirely to the verification of (3.9).

5. To establish (3.9) we will show that by choosing ρ small it is possible to find $\varepsilon > 0$ and a constant $C > 0$, $C = C(\rho, u)$ such that

$$(3.10) \quad \|P_k(\eta_\rho u)\|_r \lesssim \frac{C}{2^{(\sigma+\varepsilon)k}} \quad \text{for all } k \geq 1.$$

Clearly it is enough to prove (3.10) only for large k . We will always assume that, say, $k \geq 100$. To economise on notations we set in the rest of the proof

$$u = \eta_\rho u \quad \text{and} \quad \tilde{u} = \phi u,$$

cf. (3.4). Thus $u, \tilde{u} \in W^{\sigma, r}$ and $\text{supp} u, \tilde{u} \subset \subset \Omega$. Moreover, we can assume that

$$\|\tilde{u}\|_{W^{\sigma, r}} \leq 1,$$

and hence

$$\|u\|_{W^{\sigma, r}} \leq C_\rho,$$

As in the proof of (3.5) we start by applying the parametrix of E to (3.4). Then apply the Littlewood-Paley projection P_k and take the L^r -norm. The resulting estimate will be

$$(3.11) \quad \|P_k u\|_r \lesssim \|P_k \Phi_{-\alpha+\beta}(\eta_{2\rho} V Q u)\|_r + R,$$

where we denoted by R the remainder

$$\begin{aligned} R = & \|P_k \Phi_{-1} \tilde{u}\|_r + \|P_k \Phi_{-\alpha+\beta-1}(V Q \tilde{u})\|_r \\ & + \|P_k \Phi_{-\alpha+\beta}(V \Phi_{\gamma-1} \tilde{u})\|_r \\ & + \|P_k \Phi_{-\alpha+\beta} v\|_r \\ & + \frac{\lambda_N}{2^{Nk}}, \quad \text{with any } N > 0. \end{aligned}$$

Here $v \in L^q$ corresponds to (3.1), and the very last term corresponds to the $C_0^\infty(\Omega)$ -remainders. All operators and functions are compactly supported in Ω .

6. Now we make the first step towards proving (3.10). Namely, we show that the bound

$$(3.12) \quad R \lesssim \frac{C}{2^{(\sigma+\varepsilon)k}}$$

holds for all k with C and ε as in (3.10). Indeed, let us treat the terms of R one-by-one.

Apply (2.6) to the first term and discover that

$$\begin{aligned} \|P_k \Phi_{-1} \tilde{u}\|_r & \lesssim 2^{-k} \|\Phi_{-1} P_{k-10 \leq \cdot \leq k+10} \tilde{u}\|_r + 2^{-Nk} \\ & \lesssim 2^{(-\sigma-1)k} + 2^{-Nk} \\ & \lesssim \frac{1}{2^{(\sigma+1)k}}. \end{aligned}$$

To estimate the second term observe that Sobolev and Holder inequalities in combination with (1.4) imply

$$\|V Q \tilde{u}\|_t \lesssim 1 \quad \text{for} \quad \frac{1}{t} = \frac{1}{r} + \frac{\alpha - \beta - \sigma}{n}, \quad \frac{1}{t} > \frac{1}{r}.$$

Hence using Bernstein inequality and (2.6) we derive that

$$\begin{aligned} \|P_k \Phi_{-\alpha+\beta-1}(V Q \tilde{u})\|_r & \lesssim 2^{nk(\frac{1}{r}-\frac{1}{t})} \|P_k \Phi_{-\alpha+\beta-1}(V Q \tilde{u})\|_t \\ & \lesssim 2^{(\alpha-\beta-\sigma)k} \left(2^{(-\alpha+\beta-1)k} + 2^{-Nk} \right) \\ & \lesssim \frac{1}{2^{(\sigma+1)k}}. \end{aligned}$$

To treat the next term notice that by Holder inequality for all $\rho > 0$

$$\|\eta_{2\rho} V \Phi_{\gamma-1}(\tilde{u})\|_t \lesssim 1 \quad \text{with} \quad \frac{1}{t} = \frac{1}{r} + \frac{\alpha - \beta - \sigma - 1}{n}.$$

There are two possibilities. If $t \leq r$ then similarly to the previous argument utilise Bernstein inequality and (2.6), to discover that

$$\begin{aligned} \|P_k \Phi_{-\alpha+\beta}(V \Phi_{\gamma-1} \tilde{u})\|_r & \lesssim 2^{(\alpha-\beta-\sigma-1)k} \|P_k \Phi_{-\alpha+\beta}(V \Phi_{\gamma-1} \tilde{u})\|_t \\ & \lesssim 2^{(\alpha-\beta-\sigma-1)k} \left(2^{(-\alpha+\beta)k} + 2^{-Nk} \right) \\ & \lesssim \frac{1}{2^{(\sigma+1)k}}. \end{aligned}$$

Otherwise we have $t > r$. Then by Holder inequality

$$\|\eta_{2\rho} V \Phi_{\gamma-1} \tilde{u}\|_r \lesssim 1$$

for any small $\rho > 0$ because all functions have the compact support. We deduce from (2.6) that

$$\begin{aligned} \|P_k \Phi_{-\alpha+\beta} (V \Phi_{\gamma-1} \tilde{u})\|_r &\lesssim 2^{(-\alpha+\beta)k} + 2^{-Nk} \\ &\lesssim \frac{1}{2^{(\sigma+\varepsilon)k}}, \end{aligned}$$

where $\varepsilon = \alpha - \beta - \sigma > 0$ according to (3.6).

Finally we estimate $\|P_k \Phi_{-\alpha+\beta} v\|_r$ with $v \in L^q$. There are two possibilities. Suppose first that $q \leq r$. Then apply Bernstein inequality to discover

$$\begin{aligned} \|P_k \Phi_{-\alpha+\beta} v\|_r &\lesssim 2^{nk(\frac{\alpha-\beta-\gamma}{n} - \frac{1}{r})} \|P_k \Phi_{-\alpha+\beta} v\|_q \\ &\lesssim 2^{(\alpha-\beta-\gamma-\frac{n}{r})k} 2^{(-\alpha+\beta)k} \\ &\lesssim \frac{1}{2^{(\sigma+\varepsilon)k}}, \end{aligned}$$

where $\varepsilon = -\sigma + \gamma + n/r > 0$ due to (1.4). Alternatively, suppose $q > r$. Recall the origin of v in (3.1). We apply (2.6) and Holder inequality to deduce that

$$\begin{aligned} \|P_k \Phi_{-\alpha+\beta} v\|_r &\lesssim 2^{(-\alpha+\beta)k} \|v\|_r \\ &\lesssim \frac{1}{2^{(\sigma+\varepsilon)k}} \end{aligned}$$

with $\varepsilon = \alpha - \beta - \sigma > 0$ due to (3.6).

Combining all estimates we conclude that (3.12) holds with

$$(3.13) \quad \varepsilon = \min\{1, \alpha - \beta - \sigma, \gamma - \sigma + n/r\},$$

$\varepsilon > 0$ because of (1.4) and (3.6).

7. It is left to estimate the first term in (3.11). Applying (2.3) from Lemma 2.1 we derive that

$$\begin{aligned} \|P_k \Phi_{-\alpha+\beta} (\eta_{2\rho} V Q u)\|_r &\lesssim \|\Phi_{-\alpha+\beta} P_k (\eta_{2\rho} V Q u)\|_r \\ &\quad + \|[P_k, \Phi_{-\alpha+\beta}] u\|_r \\ &\lesssim 2^{(-\alpha+\beta)k} \|\tilde{P}_k (\eta_{2\rho} V Q u)\|_r \\ &\quad + 2^{(-\sigma-1)k} \|u\|_{W^{\sigma,r}} \\ &\lesssim \sum_{j=k-4}^{k+4} 2^{(-\alpha+\beta)j} \|P_j (\eta_{2\rho} V Q u)\|_r \\ &\quad + \frac{C_\rho}{2^{(\sigma+1)k}}. \end{aligned} \tag{3.14}$$

Thus our task is to estimate $2^{(-\alpha+\beta)k} \|P_k (\eta_{2\rho} V Q u)\|_r$. For convenience, in what follows we denote

$$V = \eta_{2\rho} V, \quad \delta = \|\eta_{2\rho} V\|_q.$$

By choosing ρ small we can make δ as small as we wish.

To estimate $P_k (V Q u)$ we follow the standard product estimates technique, see e.g. [22]. Taking into account the localisation of the Littlewood-Paley projections in the frequency space, we derive as in [14]

that for any $k \in \mathbf{Z}$

$$\begin{aligned}
P_k(VQu) &= \sum_{i,j \in \mathbf{Z}} P_k(P_i V P_j Qu) \\
&= \left\{ \sum_{i,j \in LL} + \sum_{i,j \in LH} + \sum_{i,j \in HL} + \sum_{i,j \in HH} \right\} P_k(P_i V P_j Qu) \\
&= I + II + III + IV,
\end{aligned}$$

where LL , LH , HL , and HH are the low-low, low-high, high-low, and high-high frequencies interaction zones on the integer lattice:

$$\begin{aligned}
LL &= \{i, j \in \mathbf{Z}: k-5 \leq i, j \leq k+7, \min\{i, j\} \leq k+5\}, \\
LH &= \{i, j \in \mathbf{Z}: i < k-5, k-3 \leq j \leq k+3\}, \\
HL &= \{i, j \in \mathbf{Z}: k-3 \leq i \leq k+3, j < k-5\}, \\
HH &= \{i, j \in \mathbf{Z}: i, j > k+5, |i-j| \leq 3\}.
\end{aligned}$$

We are going to estimate the four terms separately. Doing so we will constantly rely the following consequence of (2.6). For any $N \lesssim 1$ the inequality

$$(3.15) \quad \|P_j(Qu)\|_r \lesssim 2^{\gamma j} \sum_{i=j-10}^{j+10} \|P_i u\|_r + 2^{-Nj} C_\rho$$

holds for all j . As it was mentioned in (3.10) we always assume that $k \geq 100$.

8. By the properties of P_k , Bernstein inequality, and (3.15)

$$\begin{aligned}
\|I\|_r &\lesssim \sum_{i,j \in LL} \|P_i V P_j Qu\|_r \\
&\lesssim \sum_{i,j \in LL} \|P_i V\|_\infty \|P_j Qu\|_r \\
&\lesssim 2^{nk/q\delta} \sum_{j=k-5}^{k+7} \|P_j Qu\|_r \\
&\lesssim \delta 2^{k(\alpha-\beta-\gamma)} \sum_{j=k-20}^{k+20} 2^{\gamma j} \|P_j u\|_r \\
&\quad + \frac{C_\rho}{2^{Nk}}.
\end{aligned}$$

Term II is estimated exactly the same way. Thus by choosing appropriate $N \lesssim 1$ we derive

$$\begin{aligned}
2^{(-\alpha+\beta+\sigma)k} (\|I\|_r + \|II\|_r) &\lesssim \delta \sum_{j=k-20}^{k+20} 2^{\sigma j} \|P_j u\|_r \\
&\quad + \frac{C_\rho}{2^{100k}}.
\end{aligned}
\tag{3.16}$$

9. To estimate III we distinguish two cases. First, assume that

$$(3.17) \quad r \geq q.$$

Apply the Holder inequality to derive

$$\begin{aligned}
\|III\|_r &\lesssim \|(P_{k-3\leq \cdot \leq k+3} V)(P_{\cdot \leq 0} Qu)\|_r \\
&\quad + \sum_{j=1}^{k-5} \|(P_{k-3\leq \cdot \leq k+3} V)P_j Qu\|_r \\
&\lesssim \|P_{k-3\leq \cdot \leq k+3} V\|_r \|P_{\cdot \leq 0} Qu\|_\infty \\
&\quad + \sum_{j=1}^{k-5} \|P_{k-3\leq \cdot \leq k+3} V\|_r \|P_j Qu\|_\infty \\
&= X + Y.
\end{aligned}$$

From the Bernstein inequalities and (3.15) deduce that

$$\begin{aligned}
X &\lesssim 2^{nk(\frac{1}{q}-\frac{1}{r})} \|V\|_q \|P_{\cdot \leq 5} u\|_r \\
&\lesssim 2^{(\alpha-\beta-\gamma-\frac{n}{r})k} \delta C_\rho,
\end{aligned}$$

and similarly

$$\begin{aligned}
Y &\lesssim \sum_{j=1}^{k-5} 2^{(\alpha-\beta-\gamma-\frac{n}{r})k} \delta 2^{\frac{nj}{r}} \|P_j Qu\|_r \\
&\lesssim \delta \sum_{j=-10}^{k+10} 2^{(\alpha-\beta-\gamma-\frac{n}{r})k} 2^{(\frac{n}{r}+\gamma)j} \|P_j u\|_r \\
&\quad + \sum_{j=-10}^{k+10} 2^{(\alpha-\beta-\gamma-\frac{n}{r})k} \delta 2^{\frac{nj}{r}} \frac{C_\rho}{2^{Nj}}.
\end{aligned}$$

Consequently, in the case of (3.17), after a proper choice of N , we can write the final estimate for III as

$$\begin{aligned}
2^{(-\alpha+\beta+\sigma)k} \|III\|_r &\lesssim \delta \sum_{j=1}^{k+10} 2^{(\sigma-\gamma-\frac{n}{r})(k-j)} (2^{\sigma j} \|P_j u\|_r) \\
&\quad + 2^{(\sigma-\gamma-\frac{n}{r})k} C_\rho
\end{aligned} \tag{3.18}$$

Notice that $\sigma - \gamma - n/r < 0$ due to (1.4).

Next assume that

$$r < q. \tag{3.19}$$

Hence

$$\frac{1}{q} + \frac{1}{t} = \frac{1}{r} \quad \text{and} \quad q, t > r.$$

By the Holder inequality

$$\begin{aligned}
\|III\|_r &\lesssim \|P_{k-3\leq \cdot \leq k+3} V\|_q \|P_{\cdot \leq 0} (Qu)\|_t \\
&\quad + \sum_{j=1}^{k-5} \|P_{k-3\leq \cdot \leq k+3} V\|_q \|P_j Qu\|_t \\
&= Z + W.
\end{aligned}$$

The Bernstein inequalities imply that

$$Z \lesssim \delta C_\rho,$$

and

$$\begin{aligned}
W &\lesssim \delta \sum_{j=1}^{k-5} 2^{\frac{n_j}{q}} \|P_j Qu\|_r \\
&\lesssim \delta \sum_{j=-10}^{k+10} 2^{(\alpha-\beta-\gamma)j} 2^{\gamma j} \|P_j u\|_r \\
&\quad + \delta \sum_{j=-10}^{k+10} 2^{(\alpha-\beta-\gamma)j} \frac{C_\rho}{2^{Nj}}.
\end{aligned}$$

Consequently, in the case of (3.19), the final estimate for III can be written as

$$\begin{aligned}
2^{(-\alpha+\beta+\sigma)k} \|III\|_r &\lesssim \delta \sum_{j=1}^{k+10} 2^{(\sigma-\alpha+\beta)(k-j)} (2^{\sigma j} \|P_j u\|_r) \\
(3.20) \quad &\quad + 2^{(-\alpha+\beta+\sigma)k} C_\rho.
\end{aligned}$$

Notice that $-\alpha + \beta + \sigma < 0$ due to (3.6).

10. To estimate IV we also need to consider two cases. First assume that

$$(3.21) \quad r \geq q' \Leftrightarrow \frac{1}{r} + \frac{1}{q} \leq 1.$$

Then define $t \geq 1$ by writing

$$\frac{1}{t} = \frac{1}{r} + \frac{1}{q}.$$

Bernstein and Holder inequalities imply that

$$\begin{aligned}
\|P_k(P_i V P_j u)\|_r &\lesssim 2^{\frac{nk}{q}} \|P_k(P_i V P_j Qu)\|_t \\
&\lesssim 2^{(\alpha-\beta-\gamma)k} \delta \|P_j Qu\|_r.
\end{aligned}$$

Utilising (3.15) and summing over i and j in the HH region, we conclude that in the case of (3.21)

$$\begin{aligned}
2^{(-\alpha+\beta+\sigma)k} \|IV\|_r &\lesssim \delta \sum_{j=k-20}^{\infty} 2^{(\sigma-\gamma)(k-j)} (2^{\sigma j} \|P_j u\|_r) \\
&\quad + \delta \sum_{j=k-20}^{\infty} 2^{(\sigma-\gamma)k} \frac{C_\rho}{2^{Nj}} \\
&\lesssim \delta \sum_{j=k-20}^{\infty} 2^{(\sigma-\gamma)(k-j)} (2^{\sigma j} \|P_j u\|_r) \\
(3.22) \quad &\quad + \frac{C_\rho}{2^{100k}}.
\end{aligned}$$

Next assume that

$$(3.23) \quad r < q'.$$

By the Bernstein and Holder inequalities

$$\begin{aligned}
\|P_k(P_i V P_j Qu)\|_r &\lesssim 2^{nk(1-\frac{1}{r})} \|P_k(P_i V P_j Qu)\|_1 \\
&\lesssim 2^{nk(1-\frac{1}{r})} \|P_i V\|_q \|P_j Qu\|_{q'} \\
&\lesssim 2^{nk(1-\frac{1}{r})} \delta 2^{nj(\frac{1}{r}-1+\frac{1}{q})} \|P_j Qu\|_r.
\end{aligned}$$

After the summation over i and j lying in the HH zone and the application of (3.15), we discover that

$$\begin{aligned}
2^{(-\alpha+\beta+\sigma)k} \|IV\|_r &\lesssim \delta \sum_{j=k-20}^{\infty} 2^{(-\alpha+\beta+\sigma+n-\frac{n}{r})(k-j)} (2^{\sigma j} \|P_j u\|_r) \\
&\quad + \delta \sum_{j=k-20}^{\infty} 2^{(-\alpha+\beta+\sigma+n-\frac{n}{r})(k-j)} 2^{-\gamma j} \frac{C_\rho}{2^{Nj}} \\
&\lesssim \delta \sum_{j=k-20}^{\infty} 2^{(-\alpha+\beta+\sigma+n-\frac{n}{r})(k-j)} (2^{\sigma j} \|P_j u\|_r) \\
&\quad + \frac{C_\rho}{2^{100k}}
\end{aligned}
\tag{3.24}$$

provided (3.23) holds.

11. Now we can prove the desired estimate (3.10). First define θ to be the smallest parameter occurring in (3.18), (3.20), (3.22), and (3.24):

$$\begin{aligned}
0 < \theta < \min \bigg\{ &\gamma + \frac{n}{r} - \sigma, \alpha - \beta - \sigma, \sigma - \gamma, \\
&-\alpha + \beta + n \left(1 - \frac{1}{r}\right) + \sigma \bigg\}.
\end{aligned}$$

According to (1.4) and (3.6) we can find such θ . Decrease θ , if necessary, so that we also have

$$\theta < \varepsilon,$$

where ε is taken from (3.13).

We continue (3.11). The first term in the right hand side there is estimated using (3.14), (3.16), (3.18), (3.20), (3.22), and (3.24). The second term in the right hand side of (3.11) is estimated by (3.12). The resulting inequality is

$$2^{\sigma k} \|P_k u\|_r \leq C_0 \delta \sum_{j=0}^{\infty} (2^{\sigma j} \|P_j u\|_r) 2^{-\theta|j-k|} + \frac{C(\rho)}{2^{\theta k}}
\tag{3.25}$$

where the positive constant C_0 does not depend on u and ρ .

Set

$$a_k = 2^{\sigma k} \|P_k u\|_r, \quad k = 0, 1, \dots$$

We will use the following iteration lemma to bound the sequence $\{a_k\}$.

Lemma 3.1 *Let $\epsilon > 0$, let δ satisfy*

$$0 < \delta < (1 - 2^{-\epsilon})/2,$$

and let a bounded sequence $\{a_k\}$ satisfy

$$a_k \leq \frac{1}{2^{\epsilon k}} + \delta \sum_{j \geq 0} \frac{a_j}{2^{2\epsilon|k-j|}} \quad \text{for } k \geq S,$$

with some $S \geq 0$. Then

$$a_k \leq M \|\{a_k\}\|_{l^\infty} \frac{1}{2^{\epsilon k}}, \quad k = 0, 1, \dots,$$

with a constant $M \geq 0$, $M = M(\epsilon, \delta, S)$.

This elementary lemma about number sequences is proved in [14]. The proof there is a careful but straightforward iteration of the assumptions.

Going back to (3.25) we first take $\epsilon = \theta/2$. Next, find $\rho > 0$ such that we have

$$C_0\delta < (1 - 2^{-\epsilon/100})/2.$$

Then in (3.25) we can choose $S = S(u, \rho)$ such that

$$a_k \leq \frac{1}{2^{\epsilon k}} + C_0\delta \sum_{j=0}^{\infty} \frac{a_j}{2^{2\epsilon|k-j|}} \quad \text{for } k \geq S.$$

Finally, observe that

$$a_k \lesssim \|u\|_{W^{\sigma,r}} \lesssim C_\rho \quad \text{for } k = 0, 1, \dots$$

All assumptions of Lemma 3.1 now hold and we derive (3.10). ■

References

- [1] T. P. Branson, Differential operators canonically associated to a conformal structure, *Math. Scand.* **57**, (1985) 293–345.
- [2] H. Brezis, T. Kato, Remarks on the Schrodinger operator with singular complex potentials, *J. Math. Pures Appl.* **58**, (1979) 137–151.
- [3] S.-Y. A. Chang, P. C. Yang, Non-linear partial differential equations in conformal geometry. *Proceedings of the International Congress of Mathematicians Vol. I (Beijing, 2002)*, 189–207, Higher Ed. Press, Beijing, 2002.
- [4] S.-Y. A. Chang, *Non-linear elliptic equations in conformal geometry*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2004.
- [5] S.-Y. A. Chang, L. Wang, P. C. Yang, Regularity of harmonic maps, *Comm. Pure Appl. Math.* **52**, (1999) 1099–1111.
- [6] S.-Y. A. Chang, M. J. Gursky, P. C. Yang, Regularity of a fourth order nonlinear PDE with critical exponent, *Amer. J. Math.* **121**, (1999) 215–257.
- [7] Z. Djadli, E. Hebey, M. Ledoux, Paneitz-type operators and applications, *Duke Math. J.* **104**, (2000) 129–169.
- [8] C. Fefferman, C. R. Graham, Q -curvature and Poincare metrics, *Math. Res. Lett.* **9**, (2002) 139–151.
- [9] C. Fefferman, C. R. Graham, Conformal invariants, *Astérisque* 1985, 95–116.
- [10] G. P. Galdi, *An introduction to the mathematical theory of the Navier-Stokes equations. Vol. II. Nonlinear steady problems*. Springer Tracts in Natural Philosophy, 39. Springer-Verlag, New York, 1994.
- [11] C. R. Graham, M. Zworski, Scattering matrix in conformal geometry, *Invent. Math.* **152**, (2003), 89–118.
- [12] C. R. Graham, R. Jenne, L. J. Mason, and G. A. J. Sparling, Conformally invariant powers of the Laplacian. I. Existence, *J. London Math. Soc.* (2) **46** (1992) 557–565.
- [13] F. Helein, *Harmonic maps, conservation laws and moving frames*, Cambridge University Press, Cambridge, 2002.
- [14] D. A. Labutin Critical regularity for elliptic equations from Littlewood-Paley theory, *Calc. Var. Partial Differential Equations*, to appear.
- [15] Y. Y. Li, Remark on some conformally invariant integral equations: the method of moving spheres, *J. Eur. Math. Soc.* **6**, (2004) 153–180.
- [16] S. Paneitz *A quadratic conformally covariant differential operator for arbitrary pseudo-Riemannian manifold* preprint
- [17] C. D. Sogge, *Fourier integrals in classical analysis* Cambridge University Press, Cambridge, 1993.
- [18] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals* Princeton University Press, Princeton, NJ, 1993.

- [19] P. Strzelecki, A new proof of regularity of weak solutions of the H -surface equation, *Calc. Var. Partial Differential Equations*, **16**, (2003) 227–242.
- [20] M. Struwe, *Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems* Springer-Verlag, New York, 2000.
- [21] M. Struwe, *Plateau's problem and the calculus of variations* Princeton University Press, Princeton, NJ, 1988.
- [22] T. Tao *Harmonic analysis in the phase plane* UCLA lecture notes 2001, <http://www.math.ucla.edu/~tao/>
- [23] M. E. Taylor, *Pseudodifferential operators* Princeton University Press, Princeton, N.J., 1981.
- [24] N. S. Trudinger, Remarks concerning the conformal deformation of Riemannian structures on compact manifolds, *Ann. Scuola Norm. Sup. Pisa* (3) **22**, (1968) 265–274.
- [25] K. K. Uhlenbeck, J. A. Viaclovsky, Regularity of weak solutions to critical exponent variational equations, *Math. Res. Lett.* **7**, (2000) 651–656.
- [26] W. von Wahl, Die stationären Gleichungen von Navier-Stokes und semilineare elliptische Systeme. (German) *Amer. J. Math.* **100**, (1978) 1173–1184.